

UT-812
TU-544, RCNS-98-06
April 1998

Gravitino Overproduction through Moduli Decay

M. Hashimoto^{1*}, Izawa K.-I.¹, M. Yamaguchi² and T. Yanagida¹

¹*Department of Physics, University of Tokyo,
Tokyo 113-0033, Japan*

²*Department of Physics, Tohoku University,
Sendai 980-8578, Japan*

Abstract

We derive cosmological constraints on the masses of generic scalar fields which decay only through gravitationally suppressed interactions into unstable gravitinos and ordinary particles in the supersymmetric standard model. For the gravitino mass 100GeV-1TeV, the scalar masses should be larger than 100TeV to keep the success of big-bang nucleosynthesis if no late-time entropy production dilutes the gravitino density.

*Research Fellow of the Japan Society for the Promotion of Science.

1 Introduction

Superstring theories have infinitely degenerate supersymmetric vacua which are continuously connected by massless scalar fields, called moduli. These moduli fields are generally expected to acquire their masses m_φ of the order of the gravitino mass $m_{3/2}$ once supersymmetry breaking effects are included. The moduli decay into two gravitinos if the masses of moduli are larger than $2m_{3/2}$.

In this paper, we consider the decay of the moduli into two gravitinos and discuss its cosmological consequences. It is known [1, 2] that the gravitino with the mass 100GeV-1TeV decays soon after the nucleosynthesis and the decay product destroys light nuclei produced in the early universe. We see that the moduli decay tends to produce too many gravitinos to keep the success of big-bang nucleosynthesis. We derive stringent constraints on the moduli masses to avoid this disaster such as $m_\varphi \gtrsim 100\text{TeV}$.

We stress that this constraint is applicable to generic scalar fields that decay only through gravitationally suppressed interactions as long as their masses are larger than the threshold of two-gravitino decay channel.

2 The Interaction

We assume one modulus field throughout the paper, though the generalization to the case of many moduli is straightforward. We set the gravitational scale $2.4 \times 10^{18}\text{GeV}$ equal to unity.

The relevant terms in the supergravity Lagrangian [3] which describe the decay of a modulus φ into gravitinos χ is given by

$$\mathcal{L} = \epsilon^{klmn} \bar{\chi}_k \bar{\sigma}_l \frac{1}{4} (K_\varphi \partial_m \varphi - K_{\varphi^*} \partial_m \varphi^*) \chi_n - e^{\frac{K}{2}} (W^* \chi_a \sigma^{ab} \chi_b + W \bar{\chi}_a \bar{\sigma}^{ab} \bar{\chi}_b), \quad (1)$$

where W denotes the superpotential and we choose the field φ so that its vacuum expectation value vanishes: $\langle \varphi \rangle = 0$. With our definition of φ , the Kähler potential K generically contains linear terms, $K = c\varphi + c^*\varphi^* + \varphi\varphi^* + \dots$, where the coefficient c is of order one.

3 The Decay

Let us begin by discussing the decay rate of the modulus. When $m_\varphi \gg 2m_{3/2}$, the order of the decay width of φ into gravitinos is given by

$$\Gamma(\varphi \rightarrow \text{gravitinos}) \sim |c|^2 m_{3/2}^2 m_\varphi, \quad (2)$$

where c denotes the coefficient of the φ term in the Kähler potential. Here, we have used Eq.(1), the gravitino equation of motion and $\langle e^{K/2} W \rangle = m_{3/2}$.

On the other hand, the order of the decay width of φ into radiation is given by

$$\Gamma(\varphi \rightarrow \text{radiation}) \sim N m_\varphi^3, \quad (3)$$

where N is the number of the decay channels. Hence the branching ratio of the decay into gravitinos turns out to be

$$B_\chi \sim \frac{|c|^2}{N} \left(\frac{m_{3/2}}{m_\varphi} \right)^2. \quad (4)$$

The modulus φ starts damped oscillation when the Hubble scale H becomes comparable to its mass m_φ . The initial amplitude of the coherent oscillation is expected to be of order one in the Planck unit. Then the modulus density ρ_φ dominates the universe at the decay time since $\Gamma_\varphi \ll H \sim m_\varphi$. The reheat temperature after the modulus decay is given by

$$T_R \sim N_*^{-\frac{1}{4}} \sqrt{\Gamma_\varphi}, \quad (5)$$

where N_* denotes the degrees of freedom at the temperature T_R .

This implies that the gravitino number density $n_{3/2}$ produced through the modulus decay at the decay time is given by

$$\frac{n_{3/2}}{s} \sim \frac{|c|^2 N_*^{-\frac{1}{4}} m_{3/2}^2}{\sqrt{N} m_\varphi^{\frac{3}{2}}}, \quad (6)$$

where s denotes the entropy density and we have used

$$B_\chi \rho_\varphi \sim m_\varphi n_{3/2}, \quad \rho_\varphi \sim N_* T_R^4, \quad s \sim N_* T_R^3. \quad (7)$$

Namely, the modulus mass is given by

$$m_\varphi \sim \left(\frac{|c|^2 N_*^{-\frac{1}{4}} m_{3/2}^2}{\sqrt{N} y_{3/2}} \right)^{\frac{2}{3}}, \quad (8)$$

where $y_{3/2} = n_{3/2}/s$.

Gravitinos are also produced by the scattering processes of the thermal radiations after the modulus decay. The contribution to the gravitino number density is given by [2]

$$\frac{n'_{3/2}}{s} \sim 10^{-3} T_R \sim 10^{-3} N_*^{-\frac{1}{4}} \sqrt{N} m_\varphi^{\frac{3}{2}}. \quad (9)$$

4 The Bound

In the previous section, we have estimated $y_{3/2} = n_{3/2}/s$ at the decay time of the modulus φ . We may derive cosmological constraints on $y_{3/2}$ from the observation of the present universe since the estimated value $y_{3/2}$ itself yields the value of $n_{3/2}/s$ at the time of the gravitino decay.

Stringent constraints are implemented to keep the successful predictions of the big-bang nucleosynthesis provided the gravitino with the mass 100GeV-1TeV mainly decays into a photon and a photino [2]:

$$y_{3/2} \lesssim 10^{-15} - 10^{-13}. \quad (10)$$

By means of Eq.(8), we obtain

$$m_\varphi \gtrsim 100\text{TeV}. \quad (11)$$

On the other hand, the constraints due to gravitinos produced by the scattering processes of the thermal radiations read as follows:

$$m_\varphi \lesssim (10^7 - 10^9)\text{TeV}, \quad (12)$$

where we have used Eq.(9).

5 Conclusion

We have derived the constraint $100\text{TeV} \lesssim m_\varphi \lesssim 10^9\text{TeV}$ on the moduli masses m_φ in the case of the unstable gravitino² with the mass $100\text{GeV}-1\text{TeV}$. This may have obvious implications for mechanisms of the moduli stabilization. Here, the moduli may be regarded as generic scalar fields that decay only through gravitationally suppressed interactions as long as their masses are larger than the threshold of two-gravitino decay channel.

In the course of the analysis, we assumed that no entropy production has diluted the modulus and gravitino densities since the modulus density once dominated the universe. In fact, entropy production may evade the constraints. New inflation and thermal inflation are possible candidates of enough entropy production. Without such inflationary dilution, the moduli masses are severely constrained. On the other hand, if the moduli masses lie in the region $100\text{TeV} \lesssim m_\varphi \lesssim 10^9\text{TeV}$, the reheat temperature of the cosmological inflation could be very high since the density of gravitinos produced just after the inflation is diluted substantially by the decay of moduli.

Acknowledgements

The work of M.Y. was supported in part by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan No. 09640333.

²On the other hand, no stringent constraint on the moduli masses is implemented due to the moduli decay in the case of the lighter gravitino $m_{3/2} \lesssim 10\text{GeV}$ [4].

References

- [1] S. Weinberg, Phys. Rev. Lett. **48** (1982) 1303;
D.V. Nanopoulos, K.A. Olive and M. Srednicki, Phys. Lett. **B127** (1983) 30;
M.Yu. Khlopov and A.D. Linde, Phys. Lett. **B138** (1984) 265;
J. Ellis, J.E. Kim and D.V. Nanopoulos, Phys. Lett. **B145** (1984) 181;
R. Juskiewicz, J. Silk and A. Stebbins, Phys. Lett. **B158** (1985) 463;
J. Ellis, D.V. Nanopoulos and S. Sarkar, Nucl. Phys. **B259** (1985) 175;
M. Kawasaki and K. Sato, Phys. Lett. **B189** (1987) 23;
J. Ellis, G.B. Gelmini, J.L. Lopez, D.V. Nanopoulos and S. Sarkar, Nucl. Phys. **B373** (1992) 399.
- [2] M. Kawasaki and T. Moroi, Prog. Theor. Phys. **93** (1995) 879.
- [3] J. Wess and J. Bagger, *Supersymmetry and Supergravity* (Princeton University Press, 1992).
- [4] H. Pagels and J.R. Primack, Phys. Rev. Lett. **48** (1982) 223;
J. Ellis, D.V. Nanopoulos and S. Sarkar, Nucl. Phys. **B259** (1985) 175;
V.S. Berezinsky, Phys. Lett. **B261** (1991) 71;
T. Moroi, H. Murayama and M. Yamaguchi, Phys. Lett. **B303** (1993) 289.